

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

SUPERWAY
DICAMBA-M
Selective Herbicide

ACTIVE CONSTITUENTS: 340 g/L MCPA (PRESENT AS THE DIMETHYLAMINE SALT)
80 g/L DICAMBA (PRESENT AS THE DIMETHYLAMINE SALT)

GROUP	1	HERBICIDE
-------	----------	-----------

For the control of certain broadleaf weeds in winter cereals, pastures,
turf and non-crop areas.

**IMPORTANT. READ THE ATTACHED LEAFLET
BEFORE USING THIS PRODUCT.**

Contents: 1L, 2.5L, 5L, 10L, 20L, 110L, 200L, 1000L

SUPERWAY GARDEN, AG & PEST PRODUCTS PTY LTD
Shed 6, Cnr Roys Road & Moroney Place
BEERWAH QLD 4159
ABN 24 056 553 274
TEL (07) 54390355

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well-ventilated area away from children, animals, food, feedstuffs, seed and fertilisers. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and products should not be burnt.

Refillable containers: Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Harmful if absorbed by skin contact, inhaled or swallowed. Will irritate the eyes and skin. DO NOT inhale vapour. Repeated exposure may cause allergic disorders. When opening the container and preparing the spray wear cotton overalls buttoned to the neck and wrist, washable hat, elbow length PVC gloves, face shield or goggles. When using the prepared spray wear cotton overalls buttoned to the neck and wrist, washable hat and elbow length PVC gloves. If product in eyes, wash out immediately with water. If product on skin, immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each days use, wash gloves, face shield and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor, or Poisons Information Centre (Phone Australia 13 11 26).

MATERIAL SAFETY DATA SHEET

Additional information is listed on the Material Safety Data Sheet which is available from the supplier.

CONDITIONS OF SALE

The use of Superway Dicamba-M Herbicide is beyond the control of the Superway Garden, Ag and Pest Products Pty Ltd and no warranty, expressed or implied, is given by Superway Garden Ag & Pest Products Pty Ltd regarding its suitability or efficacy for any purpose for which it is used by the buyer, except the non-excludable rights under State/Territory or Federal legislation.

APVMA Approval no: 61938/1L/0407, 61938/2.5L/0407, 61938/5L/0407, 61938/10L/0407, 61938/20L/0407, 61938/110L/0407, 61938/1000L/0407

Batch No:

Date of Manufacture:

DRUM MUSTER LOGO

THIS PRODUCT IS NOT
CONSIDERED TO BE A
DANGEROUS GOOD UNDER
THE AUSTRALIAN CODE FOR
THE TRANSPORT OF
DANGEROUS GOODS BY ROAD
AND RAIL

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

SUPERWAY
DICAMBA-M
Selective Herbicide

ACTIVE CONSTITUENTS: 340 g/L MCPA (PRESENT AS THE DIMETHYLAMINE SALT)
80 g/L DICAMBA (PRESENT AS THE DIMETHYLAMINE SALT)

GROUP	1	HERBICIDE
-------	----------	-----------

For the control of certain broadleaf weeds in winter cereals, pastures,
turf and non-crop areas.

THIS LEAFLET IS PART OF THE LABEL

APVMA APPROVAL NO: 61938/0407

SUPERWAY GARDEN, AG & PEST PRODUCTS PTY LTD
Shed 6, Cnr Roys Road & Moroney Place
BEERWAH QLD 4159
ABN 24 056 553 274
TEL (07) 54390355

DIRECTIONS FOR USE:**RESTRAINTS:**

DO NOT spray when rain seems likely to occur within 4 hours.

DO NOT spray when weeds are wet with dew or rain or under stress from drought, low soil fertility, extreme cold or water logging.

DO NOT spray outside crop growth stages recommended as crop damage may result.

DO NOT apply to crops or grass pastures undersown with clover, lucerne or medics.

DO NOT apply to legume pastures.

DO NOT apply to plants suffering from stress.

Crop & Situation	Weeds	State	Rate	Critical Comments
Barley, Commercial Rye, Oats, Triticale, Wheat From early tillering stage (Zadoks 21)	Capeweed, Charlock, Common Ice-Plant, seedling Docks, Doublegee, (Threecornered Jack, Spiny Emex), Hexham Scent (Melilotus), Mustards, Soldier Thistle, Sorrel seedlings, Tares, Tree Hogweed, Turnip Weed (Rapistrum rugosum), Vetches, Ward's Weed, Wild Radish, Wild Turnip (Brassica tournefortii), Wireweed (Hogweed)	All states	1L/ha	Spray when most weeds are in the 2 to 4 leaf stage. In some instances suppression only of wireweed (hogweed), capeweed and wild radish will occur.
	Seedling Cotton Thistle, Fat-Hen	Tas only		
Barley, Commercial Rye, Oats, Triticale, Wheat From early tillering stage (Zadoks 21) until the fully tillered stage and before jointing occurs (Zadoks 30)	Cotton Thistle to bud formation	Tas only	1.7L/ha	Spray when most weeds are still in the rosette or seedling stage. N.B. Wild radish may need to be resprayed (with 2,4-D or MCPA) if future germinations occur after spraying. In some instances, suppression only of wireweed (hogweed), hoary cress and climbing buckwheat will be obtained.
	Mintweed, New Zealand Spinach	Qld, NSW, ACT only		
	Capeweed, Charlock, Chickweed, Climbing Buckwheat (Black Bindweed), Clover, Common Ice-Plant, seedling Docks, Doublegee (Threecornered Jack, Spiny Emex), Fat-Hen, Hexham Scent (Melilotus), seedling Hoary Cress, Mouse-eared Chickweed, Mustards, Narrow-leaved Plantain, Saffron Thistle, Spurry, Swinecress, Tares, Tree Hogweed, Turnip Weed (Rapistrum rugosum), Variegated Thistle, Vetch, Wild Radish, Wild Turnip (Brassica tournefortii), Wireweed (Hogweed)	Qld, NSW, ACT Tas, WA, Vic only	1.4L/ha	
Barley, Commercial Rye, Triticale From early tillering stage (Zadoks 21) until the fully tillered stage before jointing occurs (Zadoks 30)	Volunteer Peas	SA only	1L/ha plus 60mL/ha Dicamba 500	Apply when most peas are 5-7 cm high.
			1L/ha plus 120mL/ha Dicamba 500	
Oats, Wheat From early tillering stage (Zadoks 21) until the fully tillered stage and before jointing occurs (Zadoks 30)				

DIRECTIONS FOR USE (continued)

Crop & Situation	Weeds	State	Rate			Critical Comments
			Per Hectare	Per 100 Litres	Per 15L Knapsack	
Grass Pastures	Bathurst Burr, Caltrop (Yellow Vine), Chicory, Crofton Weed, Cudweed, Dandelion, Deadnettle, Fleabanes, Fumitory, Groundsel, Heliotrope, Inkweed, Mintweed, Nettles, Noogoora Burr, Pennycress, Potatoweed, Ragwort, Ripplewort, Shepherd's Purse, Smartweed, Soft Rush, Stinkwort, Thistles including Cockspur, Scotch (Spear), Slender, Star, Variegated, Wireweed (Hogweed)	All states	2.8 – 4L	190mL – 270mL	60mL	Use the higher rate on larger weeds.
	Cotton Thistle	Tas only				
	Buttercups, Hoary Cress seedlings, Horehound, Khaki Weed, Purple-Top, Veined Verbena	All states	4 - 7L	470mL	100mL	
Turf	Bindy-Eye, Capeweed, Catsear, Chickweed, Clovers, Cotula, Creeping Oxalis, Cudweeds, Dandelion, Dock, Fleabanes, Fumitory, Jo-Jo (Onehunga), Lamb's Tongue, Medics, Mustards, Narrow-leaved Plantain, Pearlwort, Pennywort, Peppergrass, Radish, Scarlet Pimpernel, Slender Thistle, Soldier Thistle, Spear Thistle, Toad Rush, Variegated Thistle, Wild Turnip, Wireweed (Hogweed).	All states	Golf Tee, Golf Green: 100mL in 15L water per 150m ² Bowling Green: 900mL- in 200L water per 1400M ² Large Turf Area: (Field, Park, Golf Fairway) 6.5L in 250-400L water per 10,000m ² (1 hectare)			Apply to actively growing weeds and moist soil. DO NOT use on Lippia, Strawberry Clover or Buffalo Grass. DO NOT mow turf for two days before or after application or fertilise within two weeks of spraying.
	Fat-Hen, Purple-Top, Red Flowered Mallow, Staggerweed, Swinecress	Qld, NSW, Tas, SA, WA				
Non-Crop Situations	Artichoke, Bindweed, Fennel, Hemlock, Thistle seedlings	All states	5.2L	350mL	80mL	A repeat spray may be necessary.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MATTER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

**WITHHOLDING PERIOD: DO NOT GRAZE OR CUT FOR STOCK FOOD 7 DAYS AFTER APPLICATION.
HARVEST: WITHHOLDING PERIOD NOT REQUIRED WHEN USED AS DIRECTED.**

GENERAL INSTRUCTIONS RESISTANT WEEDS WARNING

GROUP	I	HERBICIDE
-------	----------	-----------

Superway Dicamba-M Selective Herbicide contains members of the Benzoic acid and Phenoxy groups of herbicides. Superway Dicamba-M Selective Herbicide has the Disruptors of plant cell growth mode of action. For weed resistance management Superway Dicamba-M Selective Herbicide is a Group I Herbicide. Some naturally-occurring weed biotypes resistant to Superway Dicamba-M Selective Herbicide and other Group I herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Superway Dicamba-M Selective Herbicide or any other Group I Herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Superway Garden, Ag & Pest Products Pty Ltd accepts no liability for any losses that may result from the failure of Superway Dicamba-M Selective Herbicide to control resistant weeds.

MIXING

Add the required amount of Superway Dicamba-M Selective Herbicide directly to the tank and mix well.

APPLICATION

For Boom Application: Apply by a properly calibrated boom spray using not less than 50 litres of water per hectare unless indicated otherwise in the Critical Comments column.

For Aerial Application: Apply at least 25 litres total spray volume per hectare. The aircraft should fly as low as practical under the prevailing conditions to minimise drift.

For High Volume Application: Apply minimum 1500 litres per hectare.

For Knapsack Application: A 15-litre knapsack treats 150 square metres.

NOTE: Spray when weeds are young and actively growing. A temporary wilting may be evident in crops after application. If seasonal conditions are unusually late or dry seek further advice as damage may result from spraying under these conditions. Growers should seek advice before spraying recently released cereal varieties. There is a nil tolerance of pea seed contaminates in cereal grain and less than 100% control of Volunteer Peas could result in dockage.

COMPATIBILITY

Superway Dicamba-M Selective Herbicide is compatible with Terbutryn.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions or from spraying equipment that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

Spray equipment must be calibrated accurately prior to working on crops. Keep container sealed (airtight) when not in use. Avoid spray drift and vapour movement onto susceptible crops such as cotton, tobacco, tomatoes, vines, lupins, fruit trees and ornamentals. Minimise spray drift by using low pressure and nozzles which DO NOT give a fine droplet size. DO NOT apply this product on or near desirable trees or plants or in locations where the chemical may be washed or moved into contact with their roots. DO NOT use in high wind. All spray equipment must be thoroughly washed out after use and must not be used for spraying horticultural crops. DO NOT use at a higher rate than recommended.

PROTECTION OF LIVESTOCK, WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT use container for any other purpose. DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well-ventilated area away from children, animals, food, feedstuffs, seed and fertilisers. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500m in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and products should not be burnt.

Refillable containers: Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Harmful if absorbed by skin contact, inhaled or swallowed. Will irritate the eyes and skin. DO NOT inhale vapour. Repeated exposure may cause allergic disorders. When opening the container and preparing the spray wear cotton

overalls buttoned to the neck and wrist, washable hat, elbow length PVC gloves, face shield or goggles. When using the prepared spray wear cotton overalls buttoned to the neck and wrist, washable hat and elbow length PVC gloves. If product in eyes, wash out immediately with water. If product on skin, immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each days use, wash gloves, face shield and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor, or Poisons Information Centre (Phone Australia 13 11 26).

MATERIAL SAFETY DATA SHEET

Additional information is listed on the Material Safety Data Sheet which is available from the supplier.

CONDITIONS OF SALE

The use of Superway Dicamba-M Herbicide is beyond the control of the Superway Garden, Ag and Pest Products Pty Ltd and no warranty, expressed or implied, is given by Superway Garden Ag & Pest Products Pty Ltd regarding its suitability or efficacy for any purpose for which it is used by the buyer, except the non-excludable rights under State/Territory or Federal legislation.