

# **CAUTION**

**KEEP OUT OF REACH OF CHILDREN**

**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

# **SUPERWAY GLYPHOSATE 360 HERBICIDE**

**ACTIVE CONSTITUENT:**

**360 g/L GLYPHOSATE (present as the isopropylamine salt)**

## **WEEDKILLER**

Controls a wide range of grass and broadleaf weeds in the home garden.

CONTENTS: 250mL, 500mL, 1L

**SUPERWAY GARDEN PRODUCTS PTY LTD**

**Shed 6, Cnr Roys Road & Moroney Place**

**BEERWAH QLD 4159**

**ABN 24 056 553 274**

**TEL (07) 54390355**

## HOW TO USE:

DO NOT apply to plants suffering from stress.

DO NOT mow or cut weeds for 7 days before or after application.

DO NOT water treated plants or apply if rain is expected within 6 hours.

AREA(S)	WEEDS	RATE	HOW TO APPLY
Around buildings, driveways, fence lines, garden beds, paths	Many grass and broadleaf weeds including: <u>Annual weeds:</u> Barley, brome and ryegrasses, capeweed	Mix 10mL per 1 litre water	Spray growing plants until wet.
	<u>Perennial weeds:</u> Bamboo		Spray leaf to 2 metre tall plant.
	Blady grass, Couch	Mix 15mL per litre of water	Spray growing plant until wet at early flowering stage. Repeat treatment may be necessary.
	Johnson grass, Kikuyu grass, Nut grass	Mix 10mL per litre of water	
	Pampas grass		Spray after flowering
	Paspalum, Prairie grass, Rhodes grass, Sorrel, <u>Woody weeds:</u> Blackberry		Spray after flowering
	Lantana (NSW, Qld only)		Spray complete plant

**NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER UNLESS AUTHORISED**

## HOW TO PREPARE

Product mixes easily with clean water. Using a measuring device add the required quantity to a clean plastic, aluminium or copper container.

Do not mix more than is needed. Wash sprayer thoroughly after use.

## CAUTION

Do not allow spray to contact or drift on to plants you do not want killed.

Do not allow chemical containers or spray to get into drains, sewers, ponds or streams..

Do not plant seed or seedlings into treated area for 7 days after application.

## STORAGE AND DISPOSAL

Store in the closed original container in a cool, dry place out of the reach of children. Do not store in direct sunlight. Dispose of empty container by wrapping in paper, placing in a plastic bag and putting into the garbage.

**SAFETY DIRECTIONS**

Avoid contact with eyes and skin. Wash hands after use.

**FIRST AID**

If poisoning occurs, contact a doctor or Poisons Information Centre (PHONE: 131126).

**CONDITIONS OF SALE**

The use of Superway Glyphosate 360 Herbicide is beyond the control of the Superway Garden Products Pty Ltd and no warranty, expressed or implied, is given by Superway Garden Products Pty Ltd regarding its suitability or efficacy for any purpose for which it is used by the buyer, except the non-excludable rights under State/Territory or Federal legislation.

NRA Approval No: 46673/0702

Batch No:

Date of Manufacture:

**CAUTION**  
**KEEP OUT OF REACH OF CHILDREN**  
**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

# **SUPERWAY**

## **GLYPHOSATE 360 HERBICIDE**

**ACTIVE CONSTITUENT: 360 g/L GLYPHOSATE**  
**PRESENT AS THE ISOPROPYLAMINE SALT**

<b>GROUP</b>	<b>M</b>	<b>HERBICIDE</b>
--------------	----------	------------------

A NON-SELECTIVE FOLIAR HERBICIDE FOR THE CONTROL OF A  
WIDE RANGE OF ANNUAL AND PERENNIAL GRASSES AND  
BROADLEAFED WEEDS AS PER THE  
DIRECTIONS FOR USE TABLE

**READ THE ATTACHED BOOKLET THOROUGHLY**  
**BEFORE USING THIS PRODUCT**

CONTENTS: 1L, 2.5L, 5L, 10L, 20L, 110L, 1000L

**NRA Approval No. 46673/0702**

**SUPERWAY GARDEN PRODUCTS PTY LTD**  
**Shed 6, Cnr Roys Road & Moroney Place**  
**BEERWAH QLD 4159**  
**ABN 24 056 553 274**  
**TEL (07) 54390355**

## **PROTECTION OF CROP, NATIVE AND OTHER NON-TARGET PLANTS**

DO NOT apply under meteorological conditions or from spraying equipment which could be expected to cause spray drift onto nearby susceptible plants, adjacent crops, crop lands or pastures.

## **PROTECTION OF WILDLIFE, FISH CRUSTACEANS AND ENVIRONMENT**

DO NOT contaminate streams, rivers or waterways with the product or used container. When controlling weeds near water, refer to label direction to minimise the entry of spray into the water.

## **STORAGE AND DISPOSAL**

Store in the closed, original container in a cool, well ventilated area. Do not store for prolonged periods in direct sunlight.

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

**For refillable containers:** Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

## **SAFETY DIRECTIONS**

Product will irritate eyes and skin. Avoid contact with eyes and skin. When preparing product for use, wear elbow-length PVC gloves and face shield or goggles. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash contaminated clothing, gloves and face shield or goggles. When using controlled droplet applicator, wear protective waterproof clothing and impervious footwear.

## **FIRST AID**

If poisoning occurs, contact a doctor or Poisons Information Centre (TEL 131126).

## **MATERIAL SAFETY DATA SHEET**

For additional information refer to the Material Safety Data Sheet (MSDS) which is available from the supplier.

## **CONDITIONS OF SALE**

The use of Superway Glyphosate 360 Herbicide is beyond the control of the manufacturer, therefore no warranty, expressed or implied, is given by Superway Garden Products Pty Ltd regarding its suitability or efficacy for any purpose nor can Superway Garden Products Pty Ltd accept any responsibility for any consequence whatsoever resulting from the use of this product.

## **NOT CLASSIFIED AS DANGEROUS GOODS FOR TRANSPORT BY ROAD AND RAIL.**

**In an emergency  
dial 000  
Ask for  
Fire Brigade or  
Police**

NRA APPROVAL NO: 46673/0702

DATE OF MANUFACTURE:

BATCH NO:

**CAUTION**  
**KEEP OUT OF REACH OF CHILDREN**  
**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

# **SUPERWAY**

## **GLYPHOSATE 360 HERBICIDE**

**ACTIVE CONSTITUENT: 360 g/L GLYPHOSATE  
PRESENT AS THE ISOPROPYLAMINE SALT**

<b>GROUP</b>	<b>M</b>	<b>HERBICIDE</b>
--------------	----------	------------------

A NON-SELECTIVE FOLIAR HERBICIDE FOR THE  
CONTROL OF A WIDE RANGE OF  
ANNUAL AND PERENNIAL GRASSES AND BROADLEAFED  
WEEDS AS PER THE  
DIRECTIONS FOR USE TABLE

**READ THIS BOOKLET THOROUGHLY  
BEFORE USING THIS PRODUCT**

**NRA Approval No. 46673/0702**

**SUPERWAY GARDEN PRODUCTS PTY LTD**  
**Shed 6, Cnr Roys Road & Moroney Place**  
**BEERWAH QLD 4159**  
**ABN 24 056 553 274**  
**TEL (07) 54390355**

## DIRECTIONS FOR USE

### RESTRAINTS

DO NOT spray Superway Glyphosate 360 Herbicide if rain is likely to occur within 6 hours. Heavy rainfall within 2 hours after application may wash the chemical off the foliage and a repeat treatment may be required

DO NOT add extra surfactant or mix with other agricultural chemicals, herbicide oils or any other materials unless specifically directed on the label.

### ANNUAL WEED CONTROL - ALL STATES

Weeds Controlled	Rate	Critical Comments
Amaranth ( <i>Amaranthus spp.</i> ) Barley grass ( <i>Hordeum leporinum</i> ) Barnyard grass ( <i>Echinochloa spp.</i> ) Brome grass ( <i>Bromus spp.</i> ) Caltrop ( <i>Tribulus terrestris</i> ) Canary grass ( <i>Phalaris spp.</i> ) Capeweed ( <i>Arctotheca calendula</i> ) Cereals ( <i>volunteer</i> ) Chickweed ( <i>Stellaria media</i> ) Cobbler's Peg ( <i>Bidens pilosa</i> ) Deadnettle ( <i>Lamium amplexicaule</i> ) Double Gee ( <i>Emex australis</i> ) Fumitory ( <i>Fumaria officinalis</i> ) Ground Cherry ( <i>Physalis angulata</i> ) Lesser Swinecress ( <i>Coronopus didymus</i> ) Liverseed grass ( <i>Urochloa panicoides</i> ) Mintweed ( <i>Salvia reflexa</i> ) Paradoxa grass ( <i>Pharlaris paradoxa</i> ) Paterson's Curse ( <i>Echium plantagineum</i> ) Pigweed ( <i>Portulaca oleracea</i> ) Potato weed ( <i>Galinsoga parviflora</i> ) Rye grass ( <i>Lolium rigidum</i> ) Saffron Thistle ( <i>Carthamus lanatus</i> ) Silver grass ( <i>Vulpia spp.</i> ) Sow Thistle ( <i>Sonchus oleraceus</i> ) Spear Thistle ( <i>Cirsium vulgare</i> ) Spiny Burrgrass ( <i>Cenchrus spp.</i> ) Spurge ( <i>Euphorbia spp.</i> ) Sub Clover ( <i>Trifolium subterraneum</i> ) Thornapple ( <i>Datura spp.</i> ) Wild Mustard ( <i>Sisymbrium officinale</i> ) Wild Oats ( <i>Avena spp.</i> ) Wild Turnip ( <i>Brassica tournefortii</i> ) Winter grass ( <i>Poa annua</i> ) Variegated Thistle ( <i>Silybum marianum</i> )	<b>Boom:-</b> 2-3 L/ha  <b>Handgun:-</b> 500-700mL per 100L of water  <b>Knapsack:-</b> 75-100mL per 15L of water  Wiper Equipment and Controlled Droplet Applicators – see Application Section of this label.	<b>All Weeds</b> Spray actively growing plants The taller the weed the higher the rate. As a guide use the higher rate when weeds are higher than 15 cm.  If residual activity is required, see section titled "Compatibility". To use a residual herbicide, use the herbicides that have been recommended as being compatible in accordance with their label rates.  Use Glyphosate at rates indicated in the adjacent column.

## GENERAL USES

### For all States unless noted

SITUATION	CRITICAL COMMENTS
Agricultural areas	Superway Glyphosate 360 Herbicide may be used for the control of annual and perennial weeds as directed, in agricultural land prior to the sowing of any edible or non-edible crop, but NOT prior to transplanting seedling tomatoes.
Forests	Superway Glyphosate 360 Herbicide can be used to control weeds prior to establishing nurseries, site preparation prior to planting and amongst established trees using directed spray or shielded spray or wiper equipment. Do not allow wiper surface to contact any part of the tree. Avoid spray or spray drift contacting leaves or green bark on trees as severe injury may occur.
General weed control in Domestic areas (home garden) Commercial, Industrial and Public Service Areas, agricultural buildings and other farm situations. For specific weeds, brush and woody weeds or unwanted trees refer to the appropriate tables on this label	For the control of many grasses and broadleaf weeds  RATE: 10mL PER LITRE OF WATER  <b>Apply when weeds are actively growing. Apply to ensure complete brown and uniform wetting of foliage. Visible symptoms may take from 3-7 days to develop.</b>
Horticultural Vineyards Nuts (Including Almond, Pecan, Macadamia, Pistachio, Walnut) Citrus Stone Fruit Avocado, Banana, Blueberries, Custard Apples, Duboisia, Figs (dessert), Guava, Kiwifruit, Litchi, Mango, Monstera (fruit), Olives, Paw Paw, Persimmons, Pome fruit, Raspberries, Tea	Superway Glyphosate 360 Herbicide can be applied as a directed or shielded spray or using wiper equipment. DO NOT apply as a spray near trees or vines less than 3 years old unless they are shielded from spray and spray drift. DO NOT allow wiper to contact any part of the tree or vine. For citrus, nuts, olive, Pome fruit and vineyards - Avoid spray or spray drift contacting buds, leaves, green stems, canes or laterals, suckers, fresh wounds or fruits as severe injury may occur. Tea – apply a maximum of 4L/ha by shielded boom or directed off centre nozzle or 0.5L/100L by directed handgun or knapsack to avoid application to the crop. ALL other crops: DO NOT allow spray or spray drift to contact any part of the plant including the trunk. NOTE: Extreme care is required where split bark on Kiwi fruit and green stems of paw paw are present. May be mixed with other herbicides to provide residual control of some annual weeds. See tank mixture/ compatibility for directions.
Onions Post-plant, pre-emergence application (TAS ONLY)	For control of annual weeds and suppression of perennial weeds including rope twitch. Apply this product at 1-3L/ha post sowing and at least 7 days before crop is due to emerge. DO NOT apply to emerging onion plants as severe injury will result. Use the lower rate on small, actively growing annual weeds. Increase to the higher rate for larger annual weeds (over 15cm tall) and for suppression of perennial weeds.
Pasture	DIRECTED (SPOT) APPLICATION: This product is non selective and may damage or kill any plant in the sprayed area. Re-treatment and/or pasture improvement may be necessary to restrict seedling re-establishment. SELECTIVE APPLICATION: See WIPER EQUIPMENT BOOM APPLICATION: This product may be used to suppress existing pasture species prior to re-seeding or establishment of other crops. Where spot application (spray or wiper) is undertaken, grazing stock need not be removed. CAUTION: Certain plants may be naturally toxic to stock. Where known toxic plants are present DO NOT allow stock to graze until complete brown out has occurred.
Peanuts, Cotton, Soybeans, Sugar cane, Navy beans and chickpeas  NSW, ACT, Qld only – using selective application equipment	Using Wiper equipment apply to weeds growing between crop rows or to weeds growing at least 15 cm above the crop. Do not allow wiper to contact the crop and do not allow solution to drip form the applicator or severe crop injury will result. Refer to Wiper application section of this label. Shielded Sprayers – (cotton only) apply this product to weeds growing between cotton rows. For rates of application refer to the Weeds Controlled tables. DO not apply to crops less than 20cm high. Do not allow spray or drift to contact any part of the cotton plant as severe injury may occur.
Sugar cane Ratoon Sprayout For the control of sugar cane ratoon regrowth in NSW and Qld only	Apply to actively growing ratoons 60 – 100cm tall when growing conditions are good. Do not apply if plants are under stress from waterlogging or low moisture. Rate: 4-9L/ha. Use the higher rate for control and the lower rate for suppression or where cultivation is to follow.
Cotton Pre-Harvest For the control of Bathurst Burr, Noogoora Burr, Winter annual weeds, including Sow Thistle and the seasonal suppression of Nutgrass NSW and Qld only	DO NOT APPLY TO CROPS INTENDED FOR SEED PRODUCTION Apply alone or in tank mixtures with Dropp or Harvade. Apply when at least 60% of bolls are open and immature bolls can be easily cut with a sharp knife. Where a leafy canopy limits spray coverage, reduced weed control can be expected. For best results under these conditions, delay application until the canopy re-opens following initial conditioning treatment. Where the control of Noogoora Burr or Nutgrass is required, treatments should be applied prior to the onset of frosts. When tank mixed with defoliant, a slightly higher proportion of cotton leaf may be retained, particularly where the higher rate is used. Read and follow all tank mix products label instructions. RATE: Apply 2.5L/ha for Nutgrass control and 1.25-2.5L/ha for the other weeds. Use the lower rate on light infestations of small weeds, when the crop canopy allows adequate spray coverage of the weeds. Increase to the higher rate when the crop canopy may limit spray coverage, when treating dense infestations, or when treating larger weeds.

Non-Agricultural Areas Around buildings Commercial Domestic Industrial and Public service areas Rights of way	For residual control of annual weeds Superway Glyphosate 360 Herbicide can be tank-mixed with simazine and other residual herbicides permitted for non agricultural use. Refer to "Tank—Mixtures" section.
---	--

## PERENNIAL WEED CONTROL

WEEDS CONTROLLED	STATE	APPLICATION RATE			CRITICAL COMMENTS
		Boom L/Ha	Handgun/ Knapsack Vol/1L	Low volume Product: water	
Bamboo ( <i>Bambusa</i> spp.)	All States		10mL	-	Apply to actively growing foliage and/or regrowth which is between 1m and 2m tall. Cut stump: dilute to 1:6 ie mix 1 part Superway Glyphosate 360 Herbicide plus 6 parts water. Cut stems back to 20cm high , pour mixture down hollow stem or wet the cut.
Bent grass	Vic, Tas only	2.5L	5mL	-	Apply to actively growing plants in late spring when they have some seed-head development, but before summer moisture stress. Follow-up management is required. Full disturbance with a tyned implement should follow 10-21 days after spraying. Then follow with a summer crop and/or re-seeded pasture or crop in the following autumn.
Blady grass	Qld NSW, ACT only	9L	13mL	-	Apply to actively growing plants in the early head stage.
Bitou bush/ Bone seed <i>Chrysanthemoides monilifera</i>	Qld NSW Vic, Tas, ACT only	-	5 or 10mL	1:29 or 1:19	Plants must be actively growing. Do not spray bushes under stress. Best results will be achieved during Winter when bushes are flowering. As no residual control is provided, further treatment of germinating weed seedlings may be necessary. Use the higher rate (1:19) on bushes over 1.5m high <b>LOW VOLUME APPLICATION:</b> (eg Splatter gun gas gun): Ensure spray contacts all foliage. Use the higher rate (1:19) on bushes over 1.5m high
Blackberry <i>Rubus fruticosus</i> <i>L. agg</i>	All States	-	10-13 mL	-	Apply from January to May (flowering to leaf fall). Spray plants which are not under stress to thoroughly wet all foliage but not to cause excessive run off. Use the Higher Rate for dense, old stands over 2m high. Visible symptoms may not be fully apparent until next season. Burning (after complete brown out), pasture improvement and/or further treatment may be needed to control seedlings and regrowth. The use of CDA equipment is not recommended. TAS ONLY- Do not spray bushes bearing mature fruit.
Boxthorn, African <i>Lycium ferocissimum</i>	All States	-	7-10 mL	-	Spray to wet all foliage but not to cause excessive run off. Use the Lower Rate for young bushes and the Higher Rate for bigger mature bushes. Burning (after complete brown out), pasture improvement and/or further treatment may be needed to control seedlings and regrowth. Do not spray during hot, dry summer periods. The use of CDA equipment is not recommended.
Bracken <i>Pteridium esculentum</i>	All States	9L	15 mL	-	For boom application always add Pulse <sup>1</sup> Surfactant (200mL per 100L spray). Wiper application is recommended (see Wiper equipment). Slash bracken in previous spring/summer. Apply in March-May to completely unfurled fronds, actively growing. Visible symptoms may not be visible until next season. Follow up application is recommended Preferably associated with Pasture improvement) to achieve complete control.

Carpet Grass <i>Axonopus spp.</i>	All States	3L	5mL	-	Apply to actively growing plants when most have reached the early head stage. For Couch in SA and WA apply in October - November.
Cocksfoot <i>Dactylis glomerata</i>	All States	3L	7mL	-	
Couch <i>Cynodon dactylon</i>	All States	9L	13mL	-	
Crofton Weed	Qld, NSW, ACT only	-	5mL	-	Apply to plants with full foliage which are actively growing. Spray to wet all foliage but not to cause excessive run off. Seedling germination may have to be retreated.
Flatweed/ Cats Ear <i>(Hypochoeris radicata)</i>	All States	3L	7mL	-	Apply to fully developed rosettes at the early flower stage.
Gorse (Furze)	All States	-	10mL + Pulse <sup>1</sup> 2 mL	-	Apply to actively growing plants. May be applied all year round. Always add Pulse. Spray to wet all foliage . Burning (after complete brownout) pasture improvement and/or further treatment are recommended to control seedlings and/or regrowth.
Groundsel Bush <i>(Baccharis halimifolia)</i>	Qld, NSW & ACT only	-	7mL-10mL	1:9	Apply to actively growing plants using the higher rate for plants over 2m tall. Do not spray plants during summer drought stress conditions or during winter. Spray to wet all foliage. Further treatment and/or pasture improvement are recommended to control seedlings and/or regrowth. LOW VOLUME application: (eg Splatter gun and gas gun Use 1:9 mixture (10%) of this product : water. Apply 2 x 2mL doses per 0.5m bush height. Ensure spray contacts all foliage. Use of CDA equipment is not recommended.
Guinea Grass <i>Panicum maximum</i>	All States	9L	13mL	-	Apply to actively growing plants when most have reached the early head stage. For Wiper application see Wiper equipment.
Hawthorn <i>(Crataegus spp.)</i>	NSW, Vic, Tas, SA, WA, ACT only	-	10-13mL	1:9	Spray from flowering to leaf fall. Plants must be actively growing. Spray to thoroughly wet all foliage. Use the higher rate for plants over 2m tall. Burning (after complete brownout) pasture improvement and/or further treatment are recommended to control seedlings and/or regrowth. LOW VOLUME application: (eg Splatter gun and gas gun Use 1:9 mixture (10%) of this product : water. Apply 2 x 5mL doses per 0.5m bush height. Ensure spray contacts all foliage.
Hoary Cress <i>(Cardia draba)</i>	NSW, ACT Vic, Tas only	1.5L	5mL	-	Apply to actively growing plants which are not under stress . Best results are obtained if application is made late July-September at late rosette to flowering stage. Application may be integrated with long fallow. Cultivation may start 7 days after spraying. Wiper application may be used where sufficient stem elongation occurs (see Wiper Equipment). TAS: Add a non-ionic surfactant (50-60% ai) at 200-250mL/100L spray.
Johnson Grass <i>Sorghum halepense</i> Kangaroo Grass <i>Themeda australis</i> Kikuyu Grass <i>Pennisetum clandestinum</i>	All States	6L	10mL	-	Apply to actively growing plants when most have reached the early head stage. For wiper application on Johnson grass, off-type and volunteer sorghum, see Wiper Equipment.
Lantana ( <i>Lantana camara</i> )	Qld, NSW, ACT & NT only	-	10mL	1:9	Apply to plants with full foliage which are actively growing. Spray to thoroughly wet all foliage . Do not spray when plants are drought stressed in the summer. Pasture improvement, burning (after complete brown out) and/or retreatment of seedling regrowth may be required . Addition of Pulse (200mL/100L) may improve control. LOW VOLUME application: (eg Splatter gun and gas gun Use 1:9 mixture (10%) of this product : water. Apply 2 x 2mL doses per 0.5m bush height. Ensure spray contacts all foliage. Sprinkler Sprayer: Apply 6mL of a 1:9 solution to every square metre of treated area. Use of CDA equipment is not recommended.
Love grass (African) <i>(Eragrostis curvula)</i>	NSW, ACT Vic, WA only	6L	10mL	-	Apply to actively growing plants. To restrict seedling re-establishment pasture improvement is recommended.

<i>Luwigia peruviana</i>	All States		10mL		Apply when actively growing and at or beyond the early boom stage of growth, but before autumn colour changes occur. Thorough coverage is necessary for best control.
Mistflower ( <i>Eupatorium riparium</i> )	Qld, NSW, ACT only	-	5mL	1:9	Apply to plants with full foliage which are actively growing. Spray to thoroughly wet all foliage. Further treatment and/or pasture improvement may be necessary to restrict seedling re-establishment. Sprinkler Sprayer: Apply 3mL of a 1:9 (10%) solution to every square metre of treated area.
Nutgrass ( <i>Cyperus rotundus</i> )  Does not refer to other <i>Cyperus</i> species which may be known locally as nutgrass.	All States	6L	10mL	-	<b>NON CULTIVATED SITUATIONS:</b> Apply to actively growing plants in late Summer/autumn (Feb-April) when at least 20% have reached the head stage.
		3L + 3L	7mL + 7mL	-	<b>ARABLE LAND:</b> Make first application when at least 20% of plants have reached early head stage (about Feb). Wait for most plants to re-emerge (6-8 weeks) then make a second treatment. Follow up treatments may be necessary as further plants emerge.
Pampas grass ( <i>Cortaderia sp.</i> )	All States	-	10mL or 13mL	-	Use the higher rate for plants over 1m high. Apply during spring, summer or autumn. Plants must be actively growing. For best results ensure complete coverage of foliage and apply after flowering. Plants may first be slashed or burnt and regrowth (over 1m) may be treated.
Paragrass ( <i>Brachiara mutica</i> )	All States	9L	13mL	-	Apply to actively growing plants at the early head stage.
Paspalum <i>Paspalum dilitatum</i>		6L	10mL	-	
Pellitory	NSW, ACT only	-	10mL	-	Apply to actively growing plants prior to seeding. Repeat applications may be required to control seedlings or regrowth.
Phalaris ( <i>Phalaris aquatica</i> )	NSW, ACT, Vic, SA only	3- 6L	5mL- 10mL	-	Apply in winter-spring to actively growing plants. Use lower rate where knockdown is required eg prior to burning fire breaks. Burning can usually begin 14-21 days after spraying. Use the higher rate for longer term control.
Plantains	All States	3L	7mL	-	Apply to actively growing plants at the early head stage. Symptoms may be slow to develop.
Prairie Grass ( <i>Bromus unioloides</i> ), Qld Blue Grass ( <i>Dichanthium sericeum</i> ), Red-leg Grass ( <i>Bothriocloa ambigua</i> ), Rhodes Grass ( <i>Chloris gayana</i> )	All States	6L	10mL	-	Apply to actively growing plants at the early head stage.
Rope twitch	Vic, Tas only	6L	10mL	-	Apply in late summer-autumn to actively growing plants with foliage at least 20cm high. To ensure maximum shoot emergence the area should NOT be cultivated in the period from the preceding Winter until the time of spraying.
Sifton bush/ Chinese scrub ( <i>Cassinia arcutata</i> )	Qld, NSW, ACT only	-	10 or 13mL	1:9	Apply to actively growing plants ensuring complete coverage. Further treatment and/or pasture improvement may be necessary to restrict seedling re-establishment. Use the higher rate on bushes over 1m tall. LOW VOLUME: Apply 40mL per 0.5m height. WIPER APPLICATION: Double pass application is required. Application is best made to small (less than 1m) green bushes.
Silverleaf nightshade ( <i>Solanum elaeagnifolium</i> )	NSW, ACT only	-	20mL	-	Use ONLY under good soil moisture conditions. Apply to actively growing plants at the late flowering to berry stage. Repeat spraying will be necessary to restrict regrowth and seedling re-establishment.
Sorrel ( <i>Rumex acetosella</i> )	All States	6L	10mL	-	Apply to actively growing plants when the majority of plants are at the early bud stage.
Soursob ( <i>Oxalis pes-caprae</i> )	NSW, ACT, Vic, Tas, WA, SA only	1.5L	5mL	-	Best results are achieved if plants are treated before senescence (plant yellowing) which occurs from late July to early September. Plants should be actively growing and not under stress. If foliage area has been reduced eg due to frosting or grazing allow regrowth to about 5cm before treatment.

St John's Wort ( <i>Hypericum perforatum</i> )	All States	3L	5mL	-	Apply to actively growing plants at flowering to post flowering, procumbent stem stage (about Nov-May). To prevent seedling re-establishment, pasture improvement or re-treatment may be necessary.
Sweet Briar ( <i>Rosa rubiginosa</i> )	NSW, ACT, Vic, SA, Tas, WA only	-	15 – 20mL	1:9	Apply from late flowering to leaf fall. Plants must be actively growing. Spray to thoroughly wet all foliage. Use the Lower Rate for bushes under 1.5m tall. Burning (after complete brown out), pasture improvement and/or further treatment are recommended to control seedlings and/or regrowth. LOW VOLUME APPLICATION (eg Splatter gun or gas gun): Apply 2 x 5mL dose per 0.5m bush height. Ensure spray contacts all foliage.
Thistle – Artichoke ( <i>Cynara cardunculus</i> )	SA, Vic only	3L	5mL	-	Apply at the rosette to early head stage.
Thistle – Californian ( <i>Cirsium arvense</i> )	Vic, Tas only	6L	10mL	-	Apply to actively growing plants at the flowering stage. To ensure maximum shoot emergence the area should not be cultivated prior to spraying. Re-treatment and/or pasture improvement may be necessary to restrict seedling re-establishment.
Yorkshire Fog <i>Holcus lanatus</i>	All States	3L	7mL	-	Apply to actively growing plants at the early head stage.

## UNWANTED TREES

- FOR NSW, ACT, QLD, WA, TAS, VIC, NT ONLY

Restrains: DO NOT apply to trees under stress or to trees that are not actively growing.

METHOD	UNWANTED TREES	TREE SIZE	Rate	CRITICAL COMMENTS
Stem Injection	Flooded gum, Ghost gum, Gum topped bloodwood, Messmate stringybark, Narrowleaf Ironbark, Pink bloodwood, Poplar Box, Privet, Rhus, Spotted gum, Silverleaf Ironbark, Silvertop ash (Ironbark), Swamp Mahogany, White mahogany, Willows	Basal dia. to 25cm	Undiluted 1mL/cut	Use a specially calibrated applicator which can deliver 1 or 2mL. Make a cut at an oblique angle about 5cm deep. Ensure cut penetrates the bark to the sap steam and that product is injected as soon as possible after the cut has been made. Cuts should be 13cm apart around the tree below any branching. Remove or treat branches below the cut. For multi stemmed trees treat each stem as an individual tree.
		Basal dia. over 25cm to 60cm	Undiluted 2mL/cut	
	Camphor Laurel	Basal dia. to 25cm	Mixture 1:1 2mL/cut	
		Basal dia. over 25cm to 60cm	Undiluted 2mL/cut	
Foliar application: Low volume (Gas gun or Splatter gun)	Bullich, Marri, Jarrah	0-1.5m height	1:15 Add Pulse at 20mL/10L spray mixture	Dilute product in the recommended ratio. Calibrate splatter gun to apply 5mL of solution per dose, as a fine spray. Apply 5mL per 0.5m tree height. Ensure spray contacts all foliage.
	<i>Eucalyptus spp.</i>	0-1.5m height	1:5 Add Pulse at 20mL/10L spray mixture	
Cut stump	Jarrah, Longleaf box, Marri, Messmate Strinybark, Narrowleaf peppermint	0-10cm basal dia.	1:15	Dilute the product with water at the recommended use rate. Cut tree close to the ground and immediately wet the stump surface by swab, spray, splatter gun or brush. Remove any branches on the stump and treat the surface.
	Privet, Rhus	0-30 cm basal dia.	1:1	
Foliar application: High volume (Knapsack or handgun)	Eucalyptus species Willows	0-2.0m in height	1.0 – 1.3L per 100L  When treating Eucalyptus species add Pulse at 200mL/100L spray mixture.	Ensure all foliage is wet. Use the higher rate for trees 1 to 2 m high.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

## AQUATIC WEED CONTROL

<p><b>Aquatic Areas</b> For the control of emerged weeds in all fresh and brackish water which may be flowing, non-flowing or transient. Also for weeds on sides or banks of streams, lakes and in channels and drains.</p> <p>(See Weed Table Below)</p>	ALL STATES	DO NOT use additional surfactant/wetter, unless it is approved for use in aquatic situations.			
		<p><b>General Instructions</b></p> <ol style="list-style-type: none"> <li>Reduction in effectiveness may result if more than ¼ of the above ground portion of the weed is submerged at treatment. Submerging the treated plants following treatment may result in the spray being washed from the plant surface, thus reducing effectiveness.</li> <li>Do not apply this product within 0.5 km up-stream of a potable water intake in flowing water (i.e. river or stream etc.) or within 0.5 km of a potable water intake in a standing body of water such as a lake, pond or reservoir. Applications to moving bodies of water should be made while travelling upstream wherever possible to prevent concentration of this herbicide in water. When making any bankside applications, do not overspray more than 0.5m into open water. Avoid spraying across moving bodies of water, or where weeds do not exist.</li> <li>When spraying floating weeds, use a low volume, low pressure boom sprayer, C.D.A., or sprinkler sprayer. Do not submerge the weeds when spraying as this may wash the herbicide off the leaves.</li> <li>When emerged infestations require treatment of the total surface area of impounded water, treating the area in strips may avoid sudden impact on habitat.</li> </ol>			
Weeds Controlled	State	Rate			Critical Comments
		Boom L/ha	Handgun vol/100L	Knapsack mL/15L	
Alligator weed	All States	-	1L	150mL	Apply when actively growing, from Summer through Winter. Floating form only.
Brown beetle grass	NSW, ACT only	3L	500mL	-	Apply to actively growing plants. Do not apply to partially submerged plants.
Cumbungi ( <i>Typha spp.</i> )	All States	9L	1.3L	200mL	Apply to actively growing plants at the early head to full head stage (summer-autumn). Retreatment may be required to restrict seedling re-establishment. Wiper application is recommended (not Tasmania). See <b>Wiper Equipment</b> section.
Glyceria	Tas only	6L	1L	-	Apply to actively growing plants at the mature head stage in the late summer/autumn. Not more than ¼ of the weed should be submerged at the time of treatment.
Paragrass ( <i>Brachiara mutica</i> )	All States	9L	1.3L	200mL	Apply to actively growing plants when most have reached the early head stage.
Phragmites, Common Reed ( <i>Phragmites australis</i> )	All States	9L	1.3L	200mL	Apply to actively growing, fully developed plants approaching the early head stage. Visible symptoms of control may be slow to develop and may not be fully apparent until the next season. For wiper application see <b>Wiper Equipment</b> section.
Rushes ( <i>Juncus spp.</i> )	All States	-	-	-	Apply by wiper equipment to actively growing plants. Where there is a large proportion of dead foliage, pre-slashing is recommended. Allow adequate regrowth before treatment. Refer <b>Wiper Equipment</b> section for application instructions.
Sedge, Tall ( <i>Carex appressa</i> )	NSW, ACT, Vic, Tas only	2L or 4L	500mL or 1L	75mL or 150mL	Apply to actively growing plants in flowering to post-flowering period (Oct-Apr). Use the lower rate only if the stand has been slashed prior to treatment. Retreatment may be necessary. Visible symptoms may not be fully apparent for up to 3 months. Use of CDA equipment is not recommended.
Water Couch ( <i>Paspalum distichum</i> )	All States	9L	1.3L	200mL	Apply to actively growing plants in late summer-autumn (Feb-Mar). Do not treat after March because of the onset of winter dormancy. Full results may not be visible until the following spring. Not more than ¼ of the weed should be submerged at the time of treatment.
Water Hyacinth	All States	6L - 9L	1L - 1.3L	150-200mL	Apply when actively growing and at or beyond the early bloom stage of growth. Use the higher rate for dense infestations.
Water Lettuce	All States	-	1L - 1.3L	150-200mL	Best results are obtained from mid-summer through to winter. Use the higher rate on dense infestations.
Waterlily, Yellow	All States	6L	1L	150mL	Apply when there is a maximum emergence of floating leaves. Allow 2-3 weeks for symptoms to develop, then retreat any unaffected plants. Use low volume sprayer.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**  
**WITHHOLDING PERIOD: NOT REQUIRED WHEN USED AS DIRECTED**

## GENERAL INSTRUCTIONS

Superway Glyphosate 360 Herbicide is a non-selective, water soluble liquid herbicide effective in the control of many annual and perennial grasses and broadleaf weeds in crop areas, land preparations and non-crop areas. It is not recommended for use prior to transplanting tomato seedlings. The product is inactivated in soil and does not provide residual weed control. For residual control see tank mixtures section of this label.

Superway Glyphosate 360 Herbicide is absorbed by the plant foliage and green stems and moves through the plant from point of contact to root system. Visible effects are gradual yellowing, and wilting of the plant which advances to complete browning of growth above the ground and deterioration of underground plant parts. These effects take 3 to 7 days on annual weeds, whereas on perennial weeds it may take 2 to 3 weeks or longer depending on weather conditions following spraying.

Do not disturb treated weeds by cultivation, sowing or grazing for one day after treatment of annual weeds and 7 days for perennial weeds to ensure uptake of herbicide.

No withholding period is required for this product. However to ensure herbicide absorption, grazing of treated areas should be delayed at least one day after treatment of annual weeds and 7 days for perennial weeds. Certain plants such as soursob, St Johns Wort and Bracken, may be naturally toxic to stock. Where known toxic plants are present, grazing should be delayed until complete browning of treated plants has occurred.

**Do not** treat weeds under poor growing or dormant conditions (such as occur in drought, waterlogging, disease, insect damage or following frosts) as reduced weed control may result. Reduced efficacy may also occur when treating weeds heavily covered with dust or silt.

Delay treatment of plants wet with dew or rain if water droplets run-off when plants are disturbed.

## Resistant Weeds Warning

GROUP	M	HERBICIDE
-------	---	-----------

Superway Glyphosate 360 Herbicide is a member of the glycines group of herbicides. Superway Glyphosate 360 Herbicide has the inhibitor of EPSP synthase mode of action. For weed resistance management Superway Glyphosate 360 Herbicide is a Group M herbicide.

Some naturally occurring weed biotypes resistant to Superway Glyphosate 360 Herbicide and other Group M herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Superway Glyphosate 360 Herbicide or other Group M herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, Superway Garden Products Pty Ltd accepts no liability for any losses that may result from the failure of Superway Glyphosate 360 Herbicide to control resistant weeds.

DO NOT however assume resistance without first reviewing the method of application, timing, water quality and weather conditions.

## MIXING INSTRUCTIONS

Superway Glyphosate 360 readily mixes in with clean water. Use only clean, fresh water free from soil particles or calcium/magnesium salts (hard water). If water is acidic, use a recognised buffering agent. Spray solution should be used within a maximum time of five days to avoid loss of activity.

Do not mix, store or apply this product or spray solutions of this product in galvanised or unlined steel containers or spray tanks, since a highly flammable gas mixture may be formed.

Use stainless steel, aluminium, brass, copper, fibreglass, plastic or plastic lined containers or spray tanks. Ensure the spray tank is free from any residue or previous spray materials.

Fill spray tank with half required volume of fresh water and then add measured amount of Superway Glyphosate 360 as per directions for use rate.

Agitate well (and fill with balance of water adding surfactant near end of filling to minimise foaming with hose below surface to prevent excessive foaming).

Remove hose when desired volume is reached to avoid siphoning. When adding compatible pesticides/additive (refer TANK/MIXTURES/COMPATIBILITY) add these first in half full tank with agitation and then add Superway Glyphosate 360 with the remaining water volume and finally surfactant as above, mixing thoroughly.

DO NOT use mechanical agitators as these may cause excessive foaming.

### APPLICATION INFORMATION

Superway Glyphosate 360 Herbicide is a non-selective translocated-herbicide. Direct contact, or even drift may cause severe injury or destruction to any growing crop or other desirable plants including trees.

#### Boom Equipment

Use at spray volume of 75 to 200 L/ha. When using this product at 500mL-1.5L/ha spray volumes in the range of 25-100L/ha are preferred. Fan nozzles at pressure of 240 - 280 Kpa is recommended. Boom height must be set to ensure double overlap of nozzle patterns at the top of the weed canopy.

#### Knapsack and Handgun Equipment

Adjust equipment to deliver a fine spray pattern and ensure a complete and uniform wetting of all foliage.

For handgun equipment a D6 nozzle is recommended operating at 400-700 kPa.

Do not spray in conditions conducive to spray drift.

#### Wiper Equipment

Wiper Equipment (eg. ropewick, canvas, felt or carpet applicators) may be used to apply Superway Glyphosate 360 Herbicide on to weeds growing in oilseed crops, sugarcane, cotton, seed and pod vegetables, tree and vine crops specified in this label, pasture and non-crop areas. Avoid contact with desirable vegetation. Operate wiper equipment a minimum of 10cm above crop or pasture. Weeds should be at least 15cm above the crop or pasture at time of application. Speed of travel should be no greater than 8km/h. Best results are achieved at lower speeds and where two applications are made in opposite directions (double pass).

Where weeds are of variable height, or occur in dense infestations or clumps, some plants may not be contacted by the herbicide solution. In these cases repeat treatment may be necessary.

Mix only enough herbicide solution for immediate requirements. Do not store mixed solution for more than a few days. Flush out equipment with water after use.

RATE: Mix 1 litre of this product with 2 litres of clean water to prepare a 33% solution. This product may be used according to the above directions for suppression or control of many annual and perennial weeds. See WEEDS CONTROLLED tables for specific use recommendations.

#### Low volume equipment (eg gas gun or Splatter gun)

Apply as an even spray to cover all foliage

Refer to "Weeds Controlled" for the dilution and volume of mixture to be applied. The dilution is specified as "Low volume product:water". For example, a 1:9 mixture equals one part Superway Glyphosate 360 Herbicide plus 9 parts water.

#### Controlled Droplet Application Equipment (CDA)

- Use the following table as a guide for achieving the correct application rates using the Micron<sup>1</sup> Herbi or similar equipment. See WEEDS CONTROLLED tables for specific rates and use recommendations. For hand held equipment a walking speed of approximately 1m/sec (4km/h) is recommended.

Rate of Superway Glyphosate 360 Herbicide delivered at 1m/sec	3 L/ha	6 L/ha	9 L/ha
Mixture (by volume)			
Superway Glyphosate 360 Herbicide:			
Water	1:3	1:1	2:1
Micron Herbi, nozzle	Blue	Blue	Yellow

Do not add oils to Superway Glyphosate 360 Herbicide/water mixture, otherwise difficulty in application and reduced weed control may occur.

**CAUTION:** CDA equipment produces a fine spray pattern which is not easily visible. Ensure spray pattern or drift does not contact foliage or any other green tissue of desirable plants, since severe injury or destruction may result.

### **Aerial Equipment:**

Aerial application should only be in pasture or fallow crop situations prior to establishment of field or fodder crops or new pasture and for pre-harvest application for sorghum crops. **DO NOT** use intensive cropping situations.

Use registered rates up to a maximum of 3.2 L/ha and for Micronair and boom equipment, apply in a minimum spray volume of at least 15 L/ha. Average droplet size should be 250-350 micron diameter with a swath width of 15-17 metres.

### **Application Under Hilly Terrain:**

Increase water volume to 30-80 L/ha with droplet size 10 at least 300 micron VMD on hilly terrain.

### **Application Under Hot Conditions:**

When applying in temperatures above 25°C, increase water volume to at least 30 L/ha and droplet size to at least 300 micron VMD.

**DO NOT** apply by aircraft in temperatures above 35°C.

Avoid conditions which may result in drift such as wind over 8 km/h and inversion conditions or equipment producing droplets of 150 micron or less.

## **MAINTENANCE AND CLEANING**

Only mix, store and apply spray solutions of this product in aluminium, brass, copper, fibreglass or plastic/plastic-lined containers.

**DO NOT** use galvanised steel or unlined steel (except stainless steel) as the product or spray solution may react to produce hydrogen gas which may form a highly combustible gas mixture, which can flash or explode if ignited by open flame or other ignition source. Thoroughly flush tanks, pumps and nozzles with clean fresh water after each day's spraying. Also, if using aerial application, thoroughly wash aircraft, especially landing gear to remove herbicide residue.

## **TANK MIXTURE/COMPATIBILITY**

### **Herbicides**

This product may be mixed with the following products. Read and follow all label directions, restraints, plant back periods, withholding periods, safety directions etc for the tank mix product. Chlorsulfuron (750g/kg), Metsulfuron methyl (600 g/kg), Yield<sup>1</sup>, Stomp<sup>1</sup>, Tillmaster CT<sup>1</sup>, Logran<sup>1</sup>, LVE MCPA, Goal<sup>1</sup> CT, Flandor<sup>1</sup> 500, dicamba, 2,4-D Ester, sulfometuron methyl

### **Atrazines/Simazines**

**DO NOT** apply atrazine tank mixes for barnyard grass control.

Superway Glyphosate 360 Herbicide may be tank mixed with Atrazine Flowable or Simazine Flowable for knockdown and residual weed control.

**NOTE:** Atrazine is **NOT** approved for use in non-agricultural situations.

### **Crystalline Ammonium Sulphate**

**RATE:** 2% w/v (2kg/100L spray solution).

The addition of crystalline ammonium sulphate to Superway Glyphosate 360 Herbicide, when used to control ANNUAL weeds, MAY improve the performance of Superway Glyphosate 360 Herbicide under adverse environmental conditions such as cool, cloudy weather, and assist in minimising the antagonism in tank mixes of Superway Glyphosate 360 Herbicide and flowable triazine herbicides. Use only crystalline ammonium sulphate, NOT prilled or granulated forms. Ammonium sulphate may be corrosive to metal parts of the sprayer. Thoroughly flush tanks, pumps and nozzles with water after use.

**Insecticides** – this product is compatible with the following insecticides: dimethoate, Imidan<sup>1</sup>, Le-Mat<sup>1</sup>, chlorpyrifos, Sumithion<sup>1</sup> ULV and emulsifiable concentrates of dimethoate and fenitrothion.

Pulse<sup>1</sup> Penetrant (or equivalent): It is recommended to add an organosilicone penetrant at the rate of 200mL/100L spray solution when treating bracken (boom application), gorse, Eucalypt suckers and lantana.

### **PROTECTION OF CROP, NATIVE AND OTHER NON-TARGET PLANTS**

DO NOT apply under meteorological conditions or from spraying equipment which could be expected to cause spray drift onto nearby susceptible plants, adjacent crops, crop lands or pastures.

### **PROTECTION OF WILDLIFE, FISH CRUSTACEANS AND ENVIRONMENT**

DO NOT contaminate streams, rivers or waterways with the product or used container. When controlling weeds near water, refer to label directions to minimise entry of spray into water.

### **STORAGE AND DISPOSAL**

Store in the closed, original container in a cool, well ventilated area. Do not store for prolonged periods in direct sunlight.

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

**For refillable containers:** Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

### **SAFETY DIRECTIONS**

Product will irritate eyes and skin. Avoid contact with eyes and skin. When preparing product for use, wear elbow-length PVC gloves and face shield or goggles. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash contaminated clothing, gloves and face shield or goggles. When using controlled droplet applicator, wear protective waterproof clothing and impervious footwear.

### **FIRST AID**

If poisoning occurs, contact a doctor or Poisons Information Centre (PHONE: 131126).

### **MATERIAL SAFETY DATA SHEET**

For additional information refer to the Material Safety Data Sheet (MSDS) which is available from the supplier.

### **CONDITIONS OF SALE**

The use of Superway Glyphosate 360 Herbicide is beyond the control of the Superway Garden Products Pty Ltd and no warranty, expressed or implied, is given by Superway Garden Products Pty Ltd regarding its suitability or efficacy for any purpose for which it is used by the buyer, except the non-excludable rights under State/Territory or Federal legislation.

<sup>1</sup> Not a Superway trademark